

## Preliminary phytochemical screening of secondary metabolites from *Eucalyptus grandis*'s leaves

Jagdish Suryavanshi<sup>1</sup>, Shashank Bhatt<sup>2\*</sup>

<sup>1,2</sup>Department of Biotechnology, Rajiv Gandhi Govt. P.G. College, Mandsaur, Madhya Pradesh, India

### Abstract

*Eucalyptus grandis* is one of the useful plants which is easily available and showed various pharmacological activities. These activities depend upon the presence of secondary metabolites such as alkaloids, flavonoids, saponins tannins etc. Four solvents i.e. petroleum ether, acetone, methanol, and distilled water were used for the extraction of secondary metabolites from green leaves of *Eucalyptus grandis*. In the solvents of acetone and methanol, alkaloid glycoside, phenol, lignin, sterols and tannins were present but saponins are present only in extraction of methanol. Glycosides and tannins are present in each extraction which were petroleum ether, acetone, methanol and distilled water.

**Keywords:** secondary metabolites, *Eucalyptus grandis*, saponins

### Introduction

Ayurveda is the oldest method for the treatment of various diseases in which plant and herbs extracts are used. These plant and herbs are using for the treatment and curing of various diseases like asthma, diabetes, malaria etc. Plants are a part of life which is directly effective for each organism. Plants have two types of metabolites that are primary and secondary where Primary metabolites directly affect to plant activity while secondary metabolites are supporting part of primary metabolites. The metabolites quality and quantity indicate the effectivity of plants' part against various organisms. Thus, these types of plants are known as medicinal plant of medicines in the rural areas of developing countries (Chitme *et. al.* 2003) [6]. Many people have used medicinal plants and herbal medicines for the treatment of various diseases since ancient time (Kim, 2005, Saad *et. al.* 2017) [12]. These secondary metabolites have various pharmacological activities. *Eucalyptus* is the most important plant which belongs to Myrtaceae family and

more than 800 species distributed in different regions around the world (Hassine *et. al.* 2013) [8]. In the previous study, *Eucalyptus* leaves have shown various activities such as antioxidant, antiseptic and anti-inflammatory. Its essential oil showed herbicidal (Setia *et. al.* 2013) [24], insecticidal (Rudin, 2005; Park and Shin, 2005) [21, 18], anthelmintics (Bennet and Bryant, 1996) [4], anti-tumor (Takasaki *et. al.* 1995) [26] and anti-leech (Kirton, 2005) [11].

### Material and Methods

#### Collection of Plant Material

The green leaves of *Eucalyptus grandis* were selected for the phytochemical screening of metabolic contents. These parts were collected from Mandsaur, District Mandsaur. Mandsaur forms the northern projection of Madhya Pradesh. It lies between the parallels of latitude 23° 45' 50" North and 25° 2' 55" North, and between the meridians of longitude 74° 42' 30" East and 75° 50' 20" East.




Fig 1: Map of Madhya Pradesh

### Identification of Plant and Preliminary Screening of Secondary Metabolites

The green leaves of *Eucalyptus grandis* were collected from Mandsaur Madhya Pradesh and the plant was identified by Dr. S.N. Mishra, Principal Scientist, All India Coordinated Research Project on Medicinal and Aromatic Plants, College of Horticulture, Mandsaur, affiliated to R.V.S.K.V. Vishwavidyalaya, Gwalior (M.P.). The green leaves of *Eucalyptus grandis* were crushed using mixer grinder, and kept to cold percolation process for 48 hours with petroleum ether, acetone, methanol and distilled water. After it, the extracts were filtered and used for preliminary phytochemical screening (Shashank Bhatt *et. al.*, 2012)<sup>[25]</sup>.

### Preliminary Screening of Secondary Metabolites

After the cold percolation process of *Eucalyptus grandis*, the extracts were filtered and used for preliminary phytochemical screening such as alkaloids (Iodine, Dragendorff's test and Wagner test), flavonoids (Pew's, Shinoda and NaOH tests), glycosides (Keller-Killani, Conc. H<sub>2</sub>SO<sub>4</sub>, and Molisch tests), lignin (Labat and Lignin tests), phenols (Ellagic acid and Phenol tests), saponins (Foam and Haemolysis test), sterols (Salkowski test), tannins (Gelatin tests) were carried out (Shashank Bhatt *et. al.*, 2012)<sup>[25]</sup>.

### Preliminary Screening of Phytochemical Test Phytochemical Screening

The filtrate obtained was subjected to preliminary phytochemical screening.

#### Test for Alkaloids

**Iodine Test:** Took 3 ml of test solution into test tube and added few drops of dilute iodine solution. Blue colour appeared; but disappeared on boiling and reappeared on cooling (Khandewal K.R., 2008).

**Wagner's Test:** Took 2 to 3 ml extract solution in test tube and added few drops of Wagner's reagent. Formation of reddish-brown precipitate into solution that indicates the presence of alkaloids (Kokate C. K. *et.al*; 2001)<sup>[13]</sup>.

**Dragendorff's Tests:** Took 2 to 3 ml extract in test tube and added few drops of Dragendorff's reagent. Orange brown precipitate produced that indicates the presence of alkaloids (Kokate C. K. *et.al*; 2001)<sup>[13]</sup>.

#### Test for Flavonoids

**Pew's Tests:** Zinc powder was added into 2-3 ml. extract, followed by drop wise addition of con. HCl. Formation of purple red or cherry colour indicates the presence of flavonoids (Peach K., Tracey MV. 1956)<sup>[19]</sup>.

**Shinoda Tests:-** Took 2-3 ml. sample extract in test tube and added few fragments of magnesium metal into it, followed by dropwise addition of conc. HCl. Formation of magenta colour indicates the presence of flavonoids (Kokate C. K. *et.al*; 2001)<sup>[13]</sup>.

**NaOH Tests:** Took 2-3 ml. of ample extract and few drops of sodium hydroxide solution were added into a test tube. Formation of intense yellow colour that became colourless on addition of few drops of dilute HCl indicates the presence of flavonoids (Khandewal K.R., 2008).

#### Test for Glycosides

**Keller-Killani Test:** Glacial acetic acid was added into 2 ml. sample extract and one drop 5% FeCl<sub>3</sub> and conc. H<sub>2</sub>SO<sub>4</sub>. Reddish brown color appears at the junction of the two liquid layers and the upper layer of bluish green indicates the presence of glycosides (Kokate C. K. *et.al*; 2001)<sup>[13]</sup>.

**Glycosides test:** 1 ml water was added into small amount of sample extract and shaken well. Then aqueous solution of NaOH was added. The appearance of yellow colour indicates the presence of glycosides (Treare GE, Evans WC. 1985)<sup>[27]</sup>.

**Concentrate H<sub>2</sub>SO<sub>4</sub> Test:** 2ml. glacial acetic acid, one drop of 5% FeCl<sub>3</sub> and conc. H<sub>2</sub>SO<sub>4</sub> were added into 5ml sample extract, the appearance of brown ring indicates the presence of glycosides (Khandewal K.R., 2008).

**Molisch's Test:** 2 drops of Molisch's reagent was added into 1 ml sample extract, and 2 ml of concentrate H<sub>2</sub>SO<sub>4</sub> was added into above solution. Formation of violet ring at the junction indicates the presence of glycosides (Kokate C. K. *et. al*; 2001)<sup>[13]</sup>.

#### Test for Phenols

**Ellagic Acid Test:** Took 2 ml sample extract was treated with few drops of 5% (w/v) glacial acetic acid and 5% (w/v) NaNO<sub>2</sub> solution. The solution turned muddy or niger brown precipitate occurred in the extract. It indicates the presence of phenols solution (Gibbs R.D., 1974)<sup>[7]</sup>.

**Phenol Tests:** 0.5 ml of FeCl<sub>3</sub> (w/v) solution was added into 2 ml of test solution, formation of an intense colour that indicates the presence of phenols (Gibbs R.D., 1974)<sup>[7]</sup>.

#### Test for Lignins

**Lignin test:** 2 ml of 2% (w/v) furfuraldehyde was added into the test solution. Formation of red colour indicates the presence of lignin (Gibbs R.D., 1974)<sup>[7]</sup>.

**Labat test:** 2 ml test solution was mixed with gallic acid; it developed olive green colour that indicates the positive reaction for lignins (Gibbs R.D., 1974)<sup>[7]</sup>.

#### Test for saponins

**Foam Test:** 2 ml sample extract was diluted with 20 ml of distilled water and was shaken in a graduated cylinder for 15 minutes. A 1 cm. layer of foam, indicates the presence of saponins (Kokate C. K. *et al*; 2001)<sup>[13]</sup>.

**Haemolysis Tests:** Took one drop of extract on the glass slide and added one drop of blood. Both are mixed with wood stick and observed. Hemolytic zone appeared (Kokate C.K., 1994)<sup>[14]</sup>.

#### Test for Sterols

**Salkowski's Test:** 2ml sample extract was taken into a test tube and added 2 ml chloroform. Then conc. H<sub>2</sub>SO<sub>4</sub> was added into test tube side by side. The layer of red chloroform and acid shows greenish yellow fluorescence. It indicates the presence of sterols (Kokate C. K. *et.al*; 2001)<sup>[13]</sup>.

#### Test for Tannins

**Gelatin Test:** 1 ml Gelatin (gelatin dissolves in warm water

immediately) solution was added into 2 ml sample extract. Formation of white precipitate indicates the presence of tannins (Treare GE, Evans WC. 1985)<sup>[27]</sup>.

## Result and Discussion

The cold percolation method is one of the effective methods for the extraction of primary and secondary metabolites. Four solvents i.e. petroleum ether, acetone, methanol, and distilled water were used for the extraction of secondary metabolites from green leaves of *Eucalyptus grandis*. The result of phytochemical screening of *Eucalyptus grandis* leaves was mentioned in Table-1. In the solvents of acetone and methanol, alkaloid glycoside, phenol, lignin, sterols and tannins were present but saponins are present only in extraction of methanol. Glycosides and tannins are present in each extraction which were petroleum ether, acetone, methanol and distilled water. Alkaloids, flavonoids, lignins, saponins and sterols absent in the extraction of petroleum ether and distilled water while phenol was present in the extraction of distilled water. Phenol contents absent in the extract of petroleum ether. Acetone and methanol are an effective solvent that have the capacity to extraction of secondary metabolites in 48 hrs while petroleum ether and distilled water is mild solvent for extraction of secondary metabolites from green leaves of *E. grandis* plant.

Each secondary metabolic content such as alkaloids, flavonoids, saponins, tannins etc. has specific pharmacological activity such as anti-diabetic, anti-inflammatory, anti-cancerous etc. The flavonoids showed ant-allergic, antimicrobial and anticancer activity while tannins showed general antimicrobial and antioxidant activities (Rievere *et. al.*, 2009)<sup>[20]</sup>. Tannins may have potential value such as cytotoxic and antineoplastic agents (Aguinaldo *et. al.*, 2005)<sup>[2]</sup>. Saponins are highly effective against fungus (Aboada and Efuwape, 2001; Mohanta *et. al.*, 2007)<sup>[1, 17]</sup> that have different types of activity against different pathogens. Therefore, it can be used in the treatment of diseases. It is a bioactive antibacterial agent of plants (Mandal *et. al.* 2005; Manjunatha, 2006)<sup>[15, 16]</sup>. Steroids possess insecticidal and antimicrobial properties hence that are generally used in herbal medicines and cosmetic products (Callow; 1936)<sup>[5]</sup>. Phenolic compounds have shown anti-oxidative, antidiabetic, anticarcinogenic, antimutagenic and anti-inflammatory (Arts and Hollman; 2005, Scalbert *et. al.*; 2005)<sup>[3, 23]</sup>.

**Table 1:** *Eucalyptus grandis* Plant Seeds Phytochemical Analysis

Solvent	Petroleum Ether	Acetone	Methanol	Distilled Water
Alkaloids				
Iodine test	-ve	+ve	+ve	-ve
Dragendorff test	-ve	+ve	+ve	-ve
Wagner test	-ve	+ve	+ve	-ve
Hanger's test	-ve	+ve	+ve	-ve
Flavonoids				
Shinoda test	-ve	-ve	-ve	-ve
NaOH test	-ve	-ve	-ve	-ve
Glycosides				
Keller Killani test	+ve	+ve	+ve	+ve
Conc. H <sub>2</sub> SO <sub>4</sub> test	+ve	+ve	+ve	+ve
Molish test	-ve	+ve	+ve	+ve
Phenol				
Ellagic test	-ve	+ve	+ve	+ve
Phenol test	-ve	+ve	+ve	+ve
Lignins				

Lignin test	-ve	+ve	+ve	-ve
Labat test	-ve	+ve	+ve	-ve
Saponins				
Foam test	-ve	-ve	+ve	-ve
Haemolysis test	-ve	-ve	+ve	-ve
Sterols				
Salkowaski test	-ve	+ve	+ve	-ve
Liebermann-Burchard Test	-ve	+ve	+ve	-ve
Tannins				
Gelatin test	+ve	+ve	+ve	+ve

(+) - Indicate the Presence of Content (-) - Indicate the absence of Content

## Conclusion

Cold percolation method is one of most important method for the extraction of secondary metabolites from plant's parts. The green leaves of *Eucalyptus grandis* was used for the extraction of secondary metabolites alkaloid glycoside, phenol, lignin, sterols and tannins were present but saponins are present only in extraction of methanol. Glycosides and tannins are present in each extraction which were petroleum ether, acetone, methanol and distilled water. These contents have specific activity against specific diseases. In the four solvents, the methanol has good property for extraction of metabolic contents.

## Acknowledgement

Praying and dedicating my research article to Maa Saraswati, the goddess of knowledge and wisdom, I find no words to express my deepest gratitude to my parents, Mr. Krishna Kumar Bhatt and Mrs. Subhadra Bhatt whose encouragement helped me to go ahead on this bright path. I share the credit of my work to respectable elder brother, Mr. Mayank Bhatt who very often guided me in the work. The research work would have been a dream, had not my way been enlightened by my well-wishers and the above respectables. Last but not least, the Almighty God is unforgettable without whose kindness nothing could happen.

## References

1. Aboada OO, Efuwape BM. Antibacterial properties of some Nigerian species. Biol.Res. Comm, 2001, 13:183-188.
2. Aguinaldo AM, Espeso EI, Guevara BQ, Nonato MG. Phytochemistry. In: Guevara BQ (ed.) A Guidebook to Plant Screening: Phytochemical and Biological. University of Santo Tomas, Manila, Philippines. 2005.
3. Arts IC, Hollman PC. Polyphenols are disease risk in epidemiological studies. Am J Clin Nutr. 2005; 81(1):317s-325s.
4. Bennet JE, Bryant C. Novel source of anthelmintics. International Journal of Parasitology. 1996; 26:937-947.
5. Callow R.K. Steroids. Proc. Royal, Soc London Series, 1936, 157:194.
6. Chitme HR, Chandra R, Kaushik S. Studies on anti-diarrheal activity of *Caltropis gigantean* R. BR. In experimental animals. Journal of Pharmacy and Pharmaceutical Science. 2003; 7:70-75.
7. Gibbs RD. Chemotaxonomy of Flowering Plants. Vol.1, McGill Queen's University Press, Montreal and London, 1974.
8. Hassine DB, Abderrabbaba M, Cazaux S, Bouajila J.

- Contribution to the chemical and biological study of Eucalyptus leaves extracts: effect on frying oil. *J Med Bioeng* 2:242-247.
9. Khandelwal KR. *Practical Pharmacognocny*. 19<sup>th</sup> edition, Nirali Prakashan, Pune, 2008, p. 149-159.
  10. Khandelwal KR. *Practical Pharmacognocny*. Nirali Prakashan, Pune, edition: 19, 2008.
  11. Kirton LG. Laboratory and field test of the effectiveness of the lemon-eucalyptus extract, Citriodiol, as a repellent against land leeches of the genus *Haemadipsidae*. *Annals of Tropical Medicine & Parasitology*. 2005; 99:695-714.
  12. Kim HS. Do not put too much value on conventional medicines. *Journal of Ethnopharmacology*, 2005; 100:37-39.
  13. Kokate CK, Purohit AP, Gokhale SB. Carbohydrate and derived Products, drugs containing glycosides, drugs containing tannin Hs, lipids and protein alkaloids. Text book of Pharmacognosy, 7, edition, 2001, 133-166, 167- 254, 255-2 69, 272-310, 428-52 3.
  14. Kokate CK. *Practical Pharmacognosy*, 4th ed., Vallabh Prakasan, Delhi, 1994, 107-111.
  15. Mandal P, Sinha SP, Babu, Mandal NC. Antimicrobial activity of saponins from *Acacia auriculiformis*. *Fitoterapia*. 2005; 76(5):462-565
  16. Manjunnatha BK. Antibacterial activity of *Pterocarpus santalinus*. *Ind.J. Pharm. Sci.* 2006; 68(1):115-116.
  17. Mohanta TK, Patra JK, Rath SK, Pal DK, Thatoi HN. Evaluation of antimicrobial activity and phytochemical screening of oil and nuts of *Semicarpus anacardium* L.F. *Sci. Res. Essay*. 2007; 2(11):486-490.
  18. Park IK, Shin SC. Fumigant activity of plant essential oils and components from garlic (*Allium sativum*) and clove bud (*Eugenia caryophyllata*) oils against the japnese termite (*Reticulitermes speratus* Kolbe). *Journal of Agricultural and Food Chemistry*. 2005; 153:4388-4392.
  19. Peach K., Tracey MV. *Modern methods of plant analysis*. Vol. 3, Springer Verlag, Berlin, 1956.
  20. Rievere C, Van Nguyen JH, Pieters B Dejaegher, Heyden YV, Minh CV, Quetin-Leclercq J, et a. Polyphenols isolated from antiradial extracts of *Mallotus metcalfianus*. *Phytochemistry*. 2009; 70:86-94.
  21. Rudin W. Protection against insects. *Therapeutische Umschau*. 2005; 62:713-718.
  22. Saad AM, Mohammed MMD, Ghareeb MA, Ahmed WS, Farid MA. Chemical composition and antimicrobial activity of the essential oil of the leaves of *Cupressus macrocarpa* Hartweg. *Ex Gordon. J Appl Pharm Sci*. 2017; 7:207-212.
  23. Scalbert A, Manach C, Morand C, Remesy C, Jimenez L. Dietary polyphenols and the prevention of diseases. *Cri Rev Food Sci Nutr*. 2005; 45(4):287-306.
  24. Setia N, Batish DR, Singh HP, Kohli RK. Phytotoxicity colatile oil from *Eucalyptus citriodora* against some weedy species. *Journal of Environmental Biology*. 2007; 28:63-66.
  25. Shashank bhatt, Dr. Suresh Dhyani. Preliminary Phytochemical Screening of *Ailanthus excelsa* Roxb. *Int. J. Curr. Pharma. Res*. 2012; 4(1):87-89.
  26. Takasaki M, Konoshima T, Kozuka M, Tokuda H. Anti-tumor promoting activities of euglobals from *Eucalyptus* plants. *Biological and pharmaceutical bulletin*. 1995, 18:435-438.
  27. Treare GE, Evans WC. *Pharmacognosy* 17<sup>th</sup> edn, Bahiv Tinal, London, 1985, 149.